## SUPPLEMENTARY INFORMATION

## MULTIFUNCTIONAL MAGNETIC-POLYMERIC NANOPARTICLES BASED FERROFLUIDS FOR MULTI-MODAL IN VITRO CANCER TREATMENT USING THERMOTHERAPY AND CHEMOTHERAPY

Ganeshlenin Kandasamy <sup>1, 2</sup>, Atul Sudame <sup>1</sup>, Dipak Maity <sup>3, 1\*,</sup> Shalini Soni <sup>4</sup>, Kumari Sushmita <sup>5</sup>, Naga Suresh Veerapu <sup>4</sup>, Suryasarathi Bose <sup>5</sup>, and C V Tomy <sup>6</sup>

- <sup>1</sup> Department of Mechanical Engineering, Shiv Nadar University, Dadri, UP, India
- <sup>2</sup>Department of Biomedical Engineering, Vel Tech Rangarajan Dr. Sagunthala R&D Institute of Science and Technology, Chennai, TN, India
- <sup>3</sup> Department of Chemical Engineering, Institute of Chemical Technology Mumbai, IOC Campus, Bhubaneswar, OR, India.
- <sup>4</sup>Department of Life Sciences, Shiv Nadar University, Dadri, UP, India
- <sup>5</sup> Department of Materials Engineering, Indian Institute of Science, Bangalore, KA, India
- <sup>6</sup> School of Natural Sciences, Shiv Nadar University, Dadri, UP, India

\*Corresponding Author E-mail id: dipakmaity@gmail.com

**Supporting Figure Captions** 

**Fig. S1** Drug release profile of (i) free Cur, (ii) free Ver, (iii) Cur in S24 sample, (iv) Ver in S30 sample, (v) Cur in S32 sample, and (vi) Ver in S32 sample.

**Fig. S2** Cell viability plot of the HepG2 cancer cells incubated with the free drugs (**A**) Cur and (**B**) Ver at specific concentrations (i.e., 6.25, 12.5, 25, 50 and 100 μg/ml) for 24h.

## **Supporting Figures**

Fig S1




Fig S2

